

BMJ Open Waist-to-height ratio as an indicator of 'early health risk': simpler and more predictive than using a 'matrix' based on BMI and waist circumference

Margaret Ashwell,^{1,2} Sigrid Gibson³

To cite: Ashwell M, Gibson S. Waist-to-height ratio as an indicator of 'early health risk': simpler and more predictive than using a 'matrix' based on BMI and waist circumference. *BMJ Open* 2016;**6**:e010159. doi:10.1136/bmjopen-2015-010159

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2015-010159>).

Received 2 October 2015

Revised 6 January 2016

Accepted 29 January 2016



CrossMark

¹Ashwell Associates, Ashwell, Herts, UK

²Honorary Senior Visiting Fellow, City University London, UK

³Sig-Nurture Ltd, Guildford, Surrey, UK

Correspondence to

Dr Margaret Ashwell; margaret@ashwell.uk.com

ABSTRACT

Objectives: There is now good evidence that central obesity carries more health risks compared with total obesity assessed by body mass index (BMI). It has therefore been suggested that waist circumference (WC), a proxy for central obesity, should be included with BMI in a 'matrix' to categorise health risk. We wanted to compare how the adult UK population is classified using such a 'matrix' with that using another proxy for central obesity, waist-to-height ratio (WHtR), using a boundary value of 0.5. Further, we wished to compare cardiometabolic risk factors in adults with 'healthy' BMI divided according to whether they have WHtR below or above 0.5.

Setting, participants and outcome measures:

Recent data from 4 years (2008–2012) of the UK National Diet and Nutrition Survey (NDNS) (n=1453 adults) were used to cross-classify respondents on anthropometric indices. Regression was used to examine differences in levels of risk factors (triglycerides (TG), total cholesterol (TC), low-density lipoprotein (LDL), high-density lipoprotein (HDL), TC:HDL, glycated haemoglobin (HbA1c), fasting glucose, systolic (SBP) and diastolic blood pressure (DBP)) according to WHtR below and above 0.5, with adjustment for confounders (age, sex and BMI).

Results: 35% of the group who were judged to be at 'no increased risk' using the 'matrix' had WHtR ≥ 0.5 . The 'matrix' did not assign 'increased risk' to those with a 'healthy' BMI and 'high' waist circumference. However, our analysis showed that the group with 'healthy' BMI, and WHtR ≥ 0.5 , had some significantly higher cardiometabolic risk factors compared to the group with 'healthy' BMI but WHtR below 0.5.

Conclusions: Use of a simple boundary value for WHtR (0.5) identifies more people at 'early health risk' than does a more complex 'matrix' using traditional boundary values for BMI and WC. WHtR may be a simpler and more predictive indicator of the 'early health risks' associated with central obesity.

BACKGROUND

Anthropometric proxies for central obesity, as opposed to total obesity, assessed by body

Strengths and limitations of this study

- The use of the waist-to-height ratio (WHtR) addresses a current dilemma of how best to identify 'early health risk' with a very simple, low cost, anthropometric measure.
- The predictive value of WHtR is backed by systematic reviews and meta-analyses in many different populations.
- The analysis of cardiometabolic risk factors within the group with 'healthy' body mass index (BMI) shows that some of these factors are significantly increased if WHtR ≥ 0.5 , thus supporting the definition of WHtR >0.5 as an indicator of 'early health risk'.
- Very few studies have addressed the comparison of WHtR with any 'matrix' based on BMI and waist circumference for identifying cardiometabolic risk. We have only been able to compare our results directly with data from one other country. We hope this paper will act as a stimulus for further cost-effective analysis of existing data sets.

mass index (BMI), are usually associated with, and are slightly better predictors, of increased levels of health risk factors in populations of all ages.^{1–4} However, there have been some studies which showed that anthropometric indicators for total and central obesity did not, for example, differ in their predictive abilities.^{5–8}

In most of the above studies, waist circumference (WC) and waist-to-hip ratio (WHR)⁹ have been used as proxies for central obesity. Waist-to-height ratio (WHtR) is a proxy for central (visceral) adipose tissue,^{10–11} which has recently received attention as a marker of 'early health risk'.

A boundary value of WHtR 0.5 as a risk assessment tool was first suggested 20 years ago, and this translates into the simple message 'keep your waist to less than half your height'.^{12–15} This boundary value has

been used around the world, and findings in many populations have supported the premise that WHtR is a simple and effective anthropometric index to, for example, identify health risks.^{16–22} As well as its close relationship with morbidity, WHtR also has a clearer relationship with mortality compared with BMI.²³

We have previously shown that using BMI as a sole indicator of risk would mean that 10% of the whole UK population, and more than 25% of the UK population who are judged to be of ‘healthy’ weight using BMI, are ‘misclassified’ and might not be alerted to the need to take care or to take action.^{15 24}

The National Institute for Health and Care Excellence (NICE) tried to overcome this limitation of BMI by suggesting that WC is measured alongside BMI.²⁵ Public Health England has built on this suggestion to produce a comprehensive cross-classification matrix to categorise risk.²⁶ For simplicity and clarity, we will refer to this as the ‘matrix’. In recent guidance, NICE has advised ‘Think about using waist circumference, in addition to BMI, in people with a BMI less than 35 kg/m²’.²⁷ Earlier in 2015, indications were given that NICE wish to study research on WHtR for guidance due to be published in 2017.²⁸ Our aim was to assist NICE by comparing risk estimated by the ‘matrix’ with that estimated by WHtR.

METHODS

We used recent data from 4 years of the UK National Diet and Nutrition Survey (NDNS) (2008–2012).²⁹ The NDNS is the most authoritative source of quantitative information on the food habits and nutrient intake of the UK population. Jointly funded by the Department of Health in England (now Public Health England) and the Food Standards Agency, the results are used by government to develop policy and monitor trends in diet and nutrient intakes. Households were sampled from the UK Postcode Address File, with one adult and one child (18 months or older), or one child selected for inclusion.

We used the nurse weights for the sample (variable wt_Y1234), which adjust for unequal selection, non-response to the household/main food provider (MFP), and individual interviews and non-response to the nurse visit. Full details are given in appendix B of the survey documentation: http://doc.ukdataservice.ac.uk/doc/6533/mrdoc/pdf/6533_ndns_rp_yr1–4_userguide.pdf.

To derive the non-response weights, logistic regression modelling was used to generate a predicted probability for each participant that they would take part in the nurse’s interview based on their personal and household characteristics. These predicted probabilities were then used to generate a set of non-response weights; participants with a low predicted probability got a larger weight, increasing their representation in the sample.

Participants completed a detailed computer assisted personal interview to obtain background information (age, gender, ethnicity, region), and eating and lifestyle

behaviours such as smoking, dieting to lose weight, medication and supplement use. Anthropometric measurements (weight, height, WC) were taken by trained nurses. Weight (in bare feet and minimal clothes) was measured to the nearest 100 g using calibrated scales. Height was measured with a portable stadiometer with the head in horizontal Frankfort plane. WC was measured with a tape measure at the point midway between the iliac crest and the costal margin (lower rib). Fasting blood samples were obtained and two tubes for each participant were sent to Cambridge Addenbrooke’s for immediate analysis. Further detail of blood sampling and analysis are detailed in appendix O of survey documentation. Data files from 4 years (2008 to 2012) of the NDNS Rolling Programme were obtained under licence from the UK Data Archive (<http://www.esds.ac.uk>).

Classification of respondents by anthropometric indices (BMI, WC, WHtR)

Boundary values for WC within the ‘matrix’ were: low (men: <94 cm, women: <80 cm); high (men: 94–102 cm, women: 80–88 cm); very high (men: >102 cm, women: >88 cm). For BMI, underweight (<18.5 kg/m²); healthy weight (18.5–24.9 kg/m²); overweight (25–29.9 kg/m²); obese (30–39.9 kg/m²); very obese (>40 kg/m²).

The ‘matrix’ of WC and BMI categorises health risk as: ‘no increased risk’, ‘increased risk’, ‘high risk’ and ‘very high risk’. ‘No increased risk’ was assigned to healthy weight combined with low or high WC, and also to overweight combined with low WC. ‘Increased risk’ was assigned to healthy weight combined with very high WC, to overweight combined with high WC, to overweight combined with high WC, and to obese combined with low WC. ‘High risk’ was assigned to overweight combined with very high WC, and to obese combined with high WC. Very high risk was assigned to obese combined with very high WC, and also to very obese with any category of WC.

We combined the ‘matrix’ categories of ‘increased risk’ and ‘high risk’ to obtain 3 tiers with similar numbers of adults, for comparison with the 3 tiers of WHtR based on the following boundary values: ‘no increased risk’ (WHtR <0.5), ‘increased risk’ (WHtR ≥0.5 and <0.6) and ‘very high risk’ (WHtR ≥0.6).

The classification was based on all adults with data on BMI and WC (and WHtR) (n=1453). Data were weighted to take account of differential responses to the nurse’s visit.

Linear models to assess independent association of WHtR among ‘healthy’ weight adults

Regression models (GLM procedure in SPSS) were used to estimate the independent effect of WHtR on cardiovascular disease risk factors as outcome variables. WHtR was entered as a bivariate (<0.5 vs ≥0.5), and the models were first adjusted for age and sex, and then additionally for BMI. Interactions were not significant and were excluded from the final models. Numbers were as

follows: adults with BMI in the 'healthy' range ($n=490$) (333/157 for WHtR <0.5 vs ≥ 0.5). Two-thirds of these provided samples for analysis (213/114 for total cholesterol (TC), high-density lipoprotein (HDL) and low-density lipoprotein (LDL)), other numbers are shown in [tables 1](#) and [2](#).

RESULTS

Classification of participants

The 'matrix' categorised 41% of the NDNS adult population sampled as 'no increased risk', 32% as 'increased risk' or 'high risk', and 26% as 'very high risk' ([figure 1](#) and column b in [table 1](#)); values for men and women, respectively, 42%, 34%, 24%, and 40%, 30%, 28%, (not shown). WHtR categorised the adult population as 29% 'no increased risk', 44% 'increased risk' and 27% of the population as 'very high risk' (see [figure 1](#)). Values for men and women, respectively, were 24%, 48%, 28% and 34%, 40%, 26%, not shown in [figure 1](#).

Of greater importance, the cross tables analysis in [table 1](#) (column d) showed that more than one-third (35%) of the adult group who were judged to be at 'no increased risk' according to the 'matrix', had WHtR ≥ 0.5 , and might not be alerted to the need to take care or to take action (44% and 26% for men and women, respectively, not shown). On the contrary, [table 1](#) (column c) shows that only 3% of the group, who would be at 'increased risk' according to the 'matrix', were judged to be at 'no increased risk' using WHtR. Values for men and women, respectively, were 1% and 6%, not shown in [figure 1](#). [Figure 2](#) shows this cross tables analysis graphically for all adults.

Cardiometabolic risk factors of participants with 'healthy BMI' divided according to WHtR below and above 0.5

The classification analysis reported above showed that WHtR 0.5 classified more participants as being at 'early

increased' risk than the 'matrix'. However, the 'matrix' only classifies those with a combination of very high WC and overweight by BMI as being at 'increased' risk. People with 'healthy' BMI and high WC are defined as being at 'no increased' risk. On the contrary, classification on the basis of WHtR designates those with WHtR ≥ 0.5 , as carrying an 'increased risk', irrespective of BMI status.

We were therefore keen to investigate the people within the 'healthy' BMI range ($18.5\text{--}24.9\text{ kg/m}^2$), and we did this by comparing cardiometabolic risk factors in these adults according to their WHtR status.

[Table 2](#) shows the characteristics of the adults with BMI in the 'healthy' range when they were divided, according to WHtR, below and above 0.5. Those with WHtR ≥ 0.5 were older than those with WHtR <0.5 ($p<0.001$), and had higher mean BMI ($p<0.001$). Hence, we adjusted for age and BMI (and also for sex) in regression models.

[Table 3](#) shows that all the cardiometabolic risk factors studied were the same or indicated lower risk in the group with WHtR <0.5 . The differences for four of the risk factors, (HDL-cholesterol, TC to HDL-cholesterol ratio, triglycerides and systolic blood pressure (SBP)) reached statistical significance ($p<0.05$) when adjustment was made for age and sex. When further adjusted for BMI, as well as age and sex, three of these cardiometabolic risk factors retained statistical significance ($p<0.05$). This showed that the differences in cardiometabolic risk factors were not due to the higher BMI in the group with WHtR ≥ 0.5 .

DISCUSSION

Although BMI, WC and WHtR are, by their very nature, strongly correlated,^{24 30} the more important question is to ask which anthropometric proxy measure is the simplest and most accurate in helping to indicate 'early health risk'.

Our classification analysis showed WHtR ≥ 0.5 classified more participants as being at 'early increased' risk than the 'matrix'. Men were more likely than women to fall into this early 'increased' risk category, probably because of their greater propensity to central obesity.

We are unaware of any other UK study where risk identified by WHtR has been compared with the same 'matrix'. The New Zealand (NZ) Ministry of Health performed a similar comparison with their National Survey data.³¹ They showed that, whereas 48% of men were identified as at 'no increased risk' by the 'matrix', only 29% were classed as 'no increased risk' by WHtR <0.5 . The comparable values for women were 44% by 'matrix' and 41% by WHtR <0.5 . In other words, the NZ data also showed that WHtR 0.5 classified more people, particularly men, as being at 'early increased risk' compared with the 'matrix'.

Our analysis of the cardiometabolic risk factors in the 'healthy' BMI group made it clear that the people who




Figure 1 Categorisation of risk category of all participants by waist-to-height ratio (WHtR) and by the 'matrix'. WHtR: green = 'no increased risk', WHtR <0.5 , (29%), yellow = 'increased risk', WHtR ≥ 0.5 to <0.6 , (44%) and red = 'very high risk', WHtR $0.6+$, (27%). The 'matrix': blue = 'not applicable/underweight' (1%) green = 'no increased risk' (41%), yellow = 'increased' and 'high risk' (32%) and red = 'very high risk' (26%).

Table 1 Classification of risk category in all participants by the 'matrix', based on BMI and waist circumference, by WHtR, and by both

a	b	c	d	e
'Matrix' classification	Base (n)	WHtR <0.5 as % of those in 'matrix' category (n)	WHtR ≥0.5 and <0.6 as % in 'matrix' category (n)	WHtR ≥0.6 as % in 'matrix' category (n)
NA (underweight)	17	100% (17)	0	0
'no increased risk'	576	65% (367)	35% (209)	0
'increased' and 'high' risk	470	3% (15)	78% (366)	19% (89)
'very high risk'	390	0	18% (68)	82% (322)
Total participants	1453	29% (399)	44% (643)	27% (411)

The numbers of participants in column b are actual base numbers (unweighted). Percentages are calculated using weighted data.

The 'matrix' of waist circumference × BMI classified health risk as: 'not applicable'/underweight, 'no increased risk', 'increased risk', 'high risk' and 'very high risk'. The 'matrix' only classifies those with a combination of high waist circumference and overweight by BMI as being at 'increased' risk.

We combined the 'matrix' categories of 'increased risk' and 'high risk' to obtain three groups with similar numbers of adults, for comparison with our three categories of WHtR.

Categories for waist circumference within the 'matrix' were: low (men: <94 cm, women: <80 cm), high (men: 94–102 cm women: 80–88 cm); very high (men: >102 cm, women: >88 cm).

Categories for BMI within the 'matrix' were: underweight (<18.5 kg/m²); healthy weight (18.5–24.9 kg/m²); overweight (25–29.9 kg/m²); obese (30–39.9 kg/m²); very obese (>40 kg/m²).

Categories for WHtR were: 'no increased risk' (WHtR <0.5), 'increased risk' (WHtR 0.5 to <0.6) and 'very high risk' (WHtR 0.6+).

Columns c, d and e refer to participants in the survey who had WHtR <0.5, ≥0.5, but <0.6, and ≥0.6, expressed as a percentage of the total in each 'matrix' category.

BMI, body mass index; NA, not applicable; WHtR, waist-to-height ratio.

have a BMI in the 'healthy' range, but have WHtR >0.5, had risk factor levels that were less favourable than those in the 'healthy' BMI range with WHtR <0.5 (table 3 and figure 3). These results have prompted us to suggest that WHtR is an independent indicator of 'early health risk' after adjustment for age, sex and BMI. Given the small sample size, not all differences were statistically significant but future rounds of the NDNS rolling programme

will offer increased power. Additionally, larger data sets could be used to test the validity and generalisability of our conclusions.

There is good evidence from around the world for the metabolic implications of misclassification by BMI alone: a study of adults in Singapore found that WHtR ≥0.5 identified the highest proportion of all the cardiometabolic risk factors in men and women, even higher than a combination of BMI and WC.³² In the USA, the Bogalusa heart study of children aged from 4 to 18 years showed that nearly 10% of the children who were 'healthy' by BMI, had WHtR >0.5, and that these children had raised cardiometabolic risk factors.³³ A study using NHANES data also showed that children from 5 to 18 years with 'healthy' BMIs exhibited raised cardiometabolic risk factors if their WHtR was above 0.5.¹⁸ In Korea, the National Health and Nutrition Examination Survey showed there to be more medical concerns for




Figure 2 Percentage of participants who have waist-to-height ratio (WHtR) ≥0.5 and ≥0.6 within different categories of the 'matrix'. Vertical columns denote risk by the 'matrix': 'not applicable'; 'no increased risk', 'increased' and 'high risk' and 'very high risk'. Colours within vertical columns denote risk by WHtR: green = 'no increased risk' (WHtR <0.5), yellow = 'increased risk' (WHtR ≥0.5 to <0.6) and red = 'very high risk' (WHtR ≥0.6).

Table 2 Characteristics of adults* with 'healthy' body mass index (BMI)† according to waist-to-height ratio (WHtR) 0.5

	WHtR <0.5	WHtR ≥0.5	p Value
Number of participants*	213	114	
Sex (% men/ %women)	43/57	40/60	
Age (years) (mean, SE)	39.7 (0.9)	54.7 (1.5)	<0.001
BMI (kg/m ²) (mean, SE)	22.1 (0.09)	23.5 (0.09)	<0.001
WHtR (mean, SE)	0.46 (0.001)	0.53 (0.002)	<0.001

*With lipid values.

†BMI 'healthy' range defined as BMI 18.5–24.9 kg/m².

Table 3 Cardiometabolic risk factors for those with 'healthy' BMI* according to WHtR 0.5

	Adjusted for age and sex			Adjusted for age, sex and BMI		
	WHtR <0.5 (mean with 95% CI)	WHtR ≥0.5 (mean with 95% CI)	Difference in means	WHtR <0.5 (mean with 95% CI)	WHtR ≥0.5 (mean with 95% CI)	Difference in means
Total cholesterol (mmol/L)	4.93 (4.8 to 5.05)	4.98 (4.79 to 5.17)	0.052	4.97 (4.84 to 5.10)	4.87 (4.66 to 5.08)	-0.10
HDL-cholesterol (mmol/L)	1.67 (1.62 to 1.73)	1.49 (1.40 to 1.57)	-0.181	1.67 (1.61 to 1.73)	1.49 (1.40 to 1.59)	-0.175
Total cholesterol:	3.05 (2.93 to 3.17)	3.46 (3.28 to 3.64)	0.412	3.09 (2.97 to 3.21)	3.37 (3.18 to 3.57)	0.282
HDL-cholesterol						
Triglycerides (mmol/L)	0.99 (0.92 to 1.07)	1.15 (1.04 to 1.27)	0.162	1.02 (0.94 to 1.09)	1.09 (0.97 to 1.21)	0.072
LDL cholesterol (mmol/L)	2.84 (2.74 to 2.96)	3.00 (2.83 to 3.16)	0.148	2.88 (2.77 to 2.99)	2.91 (2.73 to 3.09)	0.029
HbA1c (%)	5.50 (5.43 to 5.58)	5.40 (5.28 to 5.51)	-0.104	5.48 (5.40 to 5.55)	5.46 (5.33 to 5.58)	-0.02
Fasting glucose (mmol/L)	5.02 (4.86 to 5.19)	4.96 (4.70 to 5.22)	-0.063	4.98 (4.81 to 5.15)	5.07 (4.82 to 4.79)	0.083
SBP (mm Hg)	121.3 (119.6 to 122.9)	124.8 (122.2 to 127.3)	3.5	121.3 (119.6 to 123.0)	124.7 (122.0 to 127.5)	3.45
DBP (mm Hg)	69.5 (68.2 to 70.7)	71.2 (69.3 to 73.1)	1.78	69.4 (68.1 to 70.6)	71.5 (69.4 to 73.5)	2.09

Estimated means were adjusted for age, sex (and BMI) as indicated.

Number of participants (n): for total cholesterol, HDL cholesterol, triglycerides=213 with WHtR <0.5 and 114 with WHtR ≥0.5; For LDL, n=213/113; HbA1c, n=208/112; glucose, n=206/105 and for SBP and DBP, n=246/119.

p Values in bold signify those which are statistically significant, that is, below 0.05.

*BMI 'healthy' range defined as BMI 18.5–24.9 kg/m².
BMI, body mass index; DBP, diastolic blood pressure; HbA1c, glycated haemoglobin; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SBP, systolic blood pressure; WHtR, waist-to-height ratio.

the adolescents in the 'healthy' weight group with central obesity (defined on the basis of WHtR) than the 'healthy' weight group without central obesity.³⁴ Further, prospective data from the ALSPAC study in UK has shown that WHtR in children aged 7–9 years predicts adolescent cardiometabolic risk better than BMI.³⁵

Our results in [table 3](#) included an analysis where the values for the cardiometabolic risk factors were adjusted for BMI as well as age and sex. Most studies, for example,^{18 24 33} do not take this 'belt and braces' approach when comparing those with WHtR below and above 0.5. However, it was reassuring that three out of four risk factors remained statistically different ($p<0.05$) even when this approach was taken. The risk factors which retained significance were HDL-cholesterol, total cholesterol: HDL-cholesterol and SBP. This gives confidence that the above-cited papers, and many others, contain valid results.

We have previously investigated adults in the NDNS²⁴ and also in the Health Survey for England,³⁶ and shown that both men and women in the group who have BMI in the 'healthy' range, but who have WHtR ≥0.5, have increased cardiometabolic risk factors, not only when they are compared with participants with 'healthy' BMI and WHtR <0.5, but even when they are compared with participants classified as overweight by BMI, who have WHtR <0.5.

The results from the cross tables analysis of our data in [table 1](#) can be extrapolated to estimate that 14% of the whole UK adult population would be 'missed' using the 'matrix'. In terms of numbers of people, and estimating the UK adult population as 48 million, this translates to nearly seven million adults who would be classified as at 'no increased risk' by the 'matrix', but




Figure 3 Participants with 'healthy' body mass index (BMI) divided according to waist-to-height ratio ≥0.5. Means for high-density lipoprotein (HDL) cholesterol and the ratio of total cholesterol: HDL cholesterol were statistically different ($p<0.05$) when adjusted for sex, age and BMI.

would have 'early increased cardiometabolic risk' identified by WHtR ≥ 0.5 .

Although our own analysis was on data from adults, many other studies suggesting the potential use of WHtR as an indicator of 'early health risk' have been performed on children or adolescents.^{33–35} There are indications that WC has increased more rapidly than BMI,³⁷ and future predictions are that this gap will widen further³⁸ reflecting the increase in central, rather than total, obesity. Ensuring that a child's WC does not exceed half his/her height can be monitored in the community and by parents. They will not even need a tape measure or weighing scales: a piece of string will suffice.^{15 39} In fact, this simple method is already recommended for self-monitoring in Thailand where the Royal Thai Ministry of Public Health has launched a campaign for adults and children to use WHtR for health promotion and prevention programmes.²¹

In terms of cost effectiveness, measuring BMI requires weighing scales as well as a stadiometer for measuring height; WHtR requires a tape measure and stadiometer. Since a tape measure is cheaper and more portable than weighing scales, we assume that the use of WHtR will be more cost effective. If the assessor only wishes to know if the participant has a WHtR at or below 0.5, the tape can be replaced by an ordinary piece of string,²¹ and the methodology becomes even more cost effective.

CONCLUSIONS AND IMPLICATIONS

WHtR is a simple primary screening risk assessment tool that identifies more people at 'early health risk' than a 'matrix', which uses a combination of BMI and WC. We recommend that the 'matrix' be amended to show that having a high WC even in the 'healthy' range of BMI, carries 'increased' risk. However, we believe that serious consideration should be given to the use of WHtR to replace the 'matrix'.

Of course, any anthropometric measure is only the first step in identifying people at 'early health risk'. More complex risk scores (eg, for diabetes) include further risk factors such as sex, age, ethnicity, socio-economic status, and family history. Further screening for clinical risk factors should follow for those deemed at risk by these simpler measures.

Our results fully support an opinion expressed very recently: '... clinicians should look beyond BMI. Although assessing for total fat mass with BMI to identify patients at greater cardiovascular risk is a good start, it is not sufficient'.⁴⁰ It is therefore timely that, in the UK, NICE intends to investigate the potential use of WHtR.²⁸

Twitter Follow Margaret Ashwell @minashwell and Sigrid Gibson @sigridgibson

Acknowledgements The authors wish to thank Dr Kirsten Rennie for additional statistical advice on the manuscript.

Contributors Both authors conceived the article and drafted the manuscript. SG analysed data from the NDNS. Both authors agreed the final manuscript.

Competing interests MA devised and copyrighted the Ashwell Shape Chart which is distributed to health professionals on a non-profit-making basis.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Janssen I, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr* 2004;79:379–84.
2. Guasch-Ferre M, Bullo M, Martinez-Gonzalez MA, *et al*. Waist-to-height ratio and cardiovascular risk factors in elderly individuals at high cardiovascular risk. *PLoS ONE* 2012;7:e43275.
3. Brambilla P, Bedogni G, Heo M, *et al*. Waist circumference-to-height ratio predicts adiposity better than body mass index in children and adolescents. *Int J Obes (Lond)* 2013;37:943–6.
4. Schneider HJ, Friedrich N, Klotsche J, *et al*. The predictive value of different measures of obesity for incident cardiovascular events and mortality. *J Clin Endocrinol Metab* 2010;95:1777–85.
5. Stevens J, Couper D, Pankow J, *et al*. Sensitivity and specificity of anthropometrics for the prediction of diabetes in a biracial cohort. *Obes Res* 2001;9:696–705.
6. Tulloch-Reid MK, Williams DE, Looker HC, *et al*. Do measures of body fat distribution provide information on the risk of type 2 diabetes in addition to measures of general obesity? Comparison of anthropometric predictors of type 2 diabetes in Pima Indians. *Diabetes Care* 2003;26:2556–61.
7. Freedman DS, Kahn HS, Mei Z, *et al*. Relation of body mass index and waist-to-height ratio to cardiovascular disease risk factors in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 2007;86:33–40.
8. Emerging Risk Factors C, Wormser D, Kaptoge S, *et al*. Separate and combined associations of body-mass index and abdominal adiposity with cardiovascular disease: collaborative analysis of 58 prospective studies. *Lancet* 2011;377:1085–95.
9. Sahakyan KR, Somers VK, Rodriguez-Escudero JP, *et al*. Normal-weight central obesity: implications for total and cardiovascular mortality. *Ann Intern Med* 2015;163:827–35.
10. Ashwell M, Cole TJ, Dixon AK. Ratio of waist circumference to height is a strong predictor of intra-abdominal fat. *BMJ* 1996;313:559–60.
11. Roriz AK, Passos LC, de Oliveira CC, *et al*. Evaluation of the accuracy of anthropometric clinical indicators of visceral fat in adults and elderly. *PLoS ONE* 2014;9:e103499.
12. Hsieh SD, Yoshinaga H. Abdominal fat distribution and coronary heart disease risk factors in men-waist/height ratio as a simple and useful predictor. *Int J Obes Relat Metab Disord* 1995;19:585–9.
13. Ashwell M, LeJeune S, McPherson K. Ratio of waist circumference to height may be a better indicator of need for weight management. *BMJ* 1996;312:377.
14. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 2005;56:303–7.
15. Ashwell M, Gibson S. A proposal for a primary screening tool: 'Keep your waist circumference to less than half your height'. *BMC Med* 2014;12:207.
16. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes Rev* 2012;13:275–86.
17. Savva SC, Lamnisos D, Kafatos AG. Predicting cardiometabolic risk: waist-to-height ratio or BMI. A meta-analysis. *Diabetes Metab Syndr Obes* 2013;6:403–19.
18. Khoury M, Manhiot C, McCrindle BW. Role of the waist/height ratio in the cardiometabolic risk assessment of children classified by body mass index. *J Am Coll Cardiol* 2013;62:742–51.
19. Jayawardana R, Ranasinghe P, Sheriff MH, *et al*. Waist to height ratio: a better anthropometric marker of diabetes and cardio-metabolic risks in South Asian adults. *Diabetes Res Clin Pract* 2013;99:292–9.

20. Rodea-Montero ER, Evia-Viscarra ML, Apolinar-Jimenez E. Waist-to-height ratio is a better anthropometric index than waist circumference and BMI in predicting metabolic syndrome among obese Mexican adolescents. *Int J Endocrinol* 2014;2014:195407.
21. Thaikruea L, Yavichai S. Proposed waist circumference measurement for waist-to-height ratio as a cardiovascular disease risk indicator: self-assessment feasibility. *Jacobs J Obes* 2015;1:1–7.
22. Liu XL, Yin FZ, Ma CP, *et al.* Waist-to-height ratio as a screening measure for identifying adolescents with hypertriglyceridemic waist phenotype. *J Pediatr Endocrinol Metab* 2015;28:1079–83.
23. Ashwell M, Mayhew L, Richardson J, *et al.* Waist-to-height ratio is more predictive of years of life lost than body mass index. *PLoS ONE* 2014;9:e103483.
24. Ashwell M, Gibson S. Waist to height ratio is a simple and effective obesity screening tool for cardiovascular risk factors: analysis of data from the British National Diet and Nutrition Survey of adults aged 19–64 years. *Obes Facts* 2009;2:97–103.
25. National Institute for Health and Clinical Excellence. NICE clinical guideline 43: Obesity: guidance on the prevention, identification, assessment and management of overweight and obesity in adults and children. London: National Institute for Health and Clinical Excellence, 2006.
26. Public Health England. Adult Obesity Slide Set. Slide 20. Adapted from: National Institute of Health and Clinical Excellence. Obesity: the prevention, identification, assessment and management of overweight and obesity in adults and children. <http://www.nice.org.uk/guidance/CG43> (accessed 18 Feb 2011). https://www.noo.org.uk/slide_sets 2013. 2015.
27. National Institute for Health and Clinical Excellence. Clinical guideline 189, Obesity: identification, assessment and management. 2014. London: National Institute for Health and Clinical Excellence, 2015.
28. Hellen N. BMI gives false hope to '7m normal weight'. *The Sunday Times* (21/6/15. 2015 21/6/15.
29. Bates B, Lennox A, Prentice A, *et al.* *National Diet and Nutrition Survey: results from Years 1–4 (combined) of the Rolling Programme (2008/2009–2011/12)*. London: Public Health England, 2014.
30. Molarius A, Seidell JC. Selection of anthropometric indicators for classification of abdominal fatness—a critical review. *Int J Obes Relat Metab Disord* 1998;22:719–27.
31. Ministry of Health. *Understanding excess body weight*. New Zealand Health Survey. Wellington: Ministry of Health, 2015.
32. Lam BC, Koh GC, Chen C, *et al.* Comparison of Body Mass Index (BMI), Body Adiposity Index (BAI), Waist Circumference (WC), Waist-To-Hip Ratio (WHR) and Waist-To-Height Ratio (WHtR) as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS ONE* 2015;10:e0122985.
33. Mokha JS, Srinivasan SR, Dasmahapatra P, *et al.* Utility of waist-to-height ratio in assessing the status of central obesity and related cardiometabolic risk profile among normal weight and overweight/obese children: the Bogalusa Heart Study. *BMC Pediatr* 2010;10:73.
34. Cho WK, Kim H, Lee HY, *et al.* Insulin Resistance of Normal Weight Central Obese Adolescents in Korea Stratified by Waist to Height Ratio: Results from the Korea National Health and Nutrition Examination Surveys 2008–2010. *Int J Endocrinol* 2015;2015:158758.
35. Graves L, Garnett SP, Cowell CT, *et al.* Waist-to-height ratio and cardiometabolic risk factors in adolescence: findings from a prospective birth cohort. *Pediatr Obes* 2014;9:327–38.
36. Gibson S, Ashwell M. Non-overweight 'apples' have higher cardiometabolic risk factors than overweight 'pears': waist-to-height ratio is a better screening tool than BMI for blood levels of cholesterol and glycated haemoglobin. *Obes Facts* 2015;8(Suppl 1):139.
37. Mindell JS, Dinsdale H, Ridler C, *et al.* Changes in waist circumference among adolescents in England from 1977–1987 to 2005–2007. *Public Health* 2012;126:695–701.
38. Shaw A, Retat L, Brown M, *et al.* Beyond BMI: projecting the future burden of obesity in England using different measures of adiposity. *Obes Facts* 2015;8(Suppl 1):137.
39. Wise J. Piece of string can assess cardiovascular risk, study finds. *BMJ* 2015;350:h2434.
40. Poirier P. The many paradoxes of our modern world: is there really an obesity paradox or is it only a matter of adiposity assessment? *Ann Intern Med* 2015;163:880–1.

BMJ Open

Waist-to-height ratio as an indicator of 'early health risk': simpler and more predictive than using a 'matrix' based on BMI and waist circumference

Margaret Ashwell and Sigrid Gibson

BMJ Open 2016 6:

doi: [10.1136/bmjopen-2015-010159](https://doi.org/10.1136/bmjopen-2015-010159)

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/6/3/e010159>

These include:

References

This article cites 34 articles, 6 of which you can access for free at:
<http://bmjopen.bmj.com/content/6/3/e010159#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Global health](#) (310)
[Health policy](#) (459)
[Public health](#) (1516)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>